

10/531,252

(FILE 'HOME' ENTERED AT 13:00:20 ON 10 NOV 2007)

FILE 'CAPLUS' ENTERED AT 13:03:48 ON 10 NOV 2007

=> s reduction of palladium

332043 REDUCTION

172232 PALLADIUM

L1 728 REDUCTION OF PALLADIUM
(REDUCTION(1W) PALLADIUM)

=> s l1 and propylene

192172 PROPYLENE

L2 4 L1 AND PROPYLENE

=> d 1-4 bib abs

L2 ANSWER 1 OF 4 CAPLUS COPYRIGHT 2007 ACS on STN

AN 2002:794351 CAPLUS

DN 137:295352

TI Method and solution-suspended particulate palladium catalyst for the manufacture of acrylic or methacrylic acids via the oxidation of propylene or isobutylene

IN Unruh, Jerry D.; Diaz, Norma Jean; Molina, Robert Ray; Snyder, Phillip Sidney; Windhorst, Kenneth Allen

PA USA

SO U.S. Pat. Appl. Publ., 10 pp.

CODEN: USXXCO

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 2002151747	A1	20021017	US 2001-833945	20010412
	CA 2412307	A1	20021024	CA 2002-2412307	20020411
	WO 2002083299	A2	20021024	WO 2002-US11386	20020411
	WO 2002083299	A3	20030227		
	W: AU, BR, CA, CN, CZ, ID, IN, JP, KR, MX, SG, US, ZA				
	RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR				
	AU 2002258767	A1	20021028	AU 2002-258767	20020411
	BR 2002004827	A	20030617	BR 2002-4827	20020411
	EP 1377372	A2	20040107	EP 2002-728734	20020411
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI, CY, TR				
	JP 2004519326	T	20040702	JP 2002-581095	20020411
	IN 2002KN01506	A	20040717	IN 2002-KN1506	20021210
	ZA 2002010038	A	20031211	ZA 2002-10038	20021211
	MX 2002PA12317	A	20040906	MX 2002-PA12317	20021211
	US 2004181082	A1	20040916	US 2003-415714	20030919
PRAI	US 2001-833945	A	20010412		
	WO 2002-US11386	W	20020411		

AB A method for the manufacture of acrylic acid or methacrylic acid by the oxidation

of propylene or isobutylene is achieved by: (a) reducing palladium acetate to unsupported palladium with propylene in an oxygen-free single- or two-phase aqueous solution containing as a co-solvent a maximum

concentration of a C2-C6 carboxylic acid or C3-C6 ketone in a reactor adapted for

continuous-phase production; (b) adding air and propylene or isobutylene in a continuous manner; (c) recovering the acrylic acid or methacrylic acid formed; and (d) recycling the solvent to the reactor.

L2 ANSWER 2 OF 4 CAPLUS COPYRIGHT 2007 ACS on STN
 AN 1995:197562 CAPLUS
 DN 122:16036
 TI The performance of Pd, Pt, and Pd-Pt catalysts in lean exhaust
 AU Beck, Donald D.; Monroe, David R.; DiMaggio, Craig L.; Sommers, John W.
 CS General Motors Research, USA
 SO Society of Automotive Engineers, [Special Publication] SP (1993),
 SP-957(U.S. and European Automotive Emissions Technology), 71-83
 CODEN: SAESA2; ISSN: 0099-5908
 DT Journal
 LA English
 AB A matrix of pelleted catalysts composed of Pt, Pd, Pt co-impregnated with
 Pd, and Pt phys. mixed with Pd supported on Al₂O₃ were compared with the
 same noble metal formulations supported on CeO₂/Al₂O₃ for light-off and
 warmed-up performance in net lean exhaust. These catalysts were tested as
 prepared (fresh) and following a relatively severe thermal aging treatment
 (cycled between net lean and net rich environment at 1000°C for 4
 h). Pd showed better light-off performance than Pt for catalyzing the
 oxidation of propylene, while Pt showed better light-off and
 warmed-up performance than Pd for catalyzing the oxidation of propane.
 Having both Pt and Pd present as a result of co-impregnation or phys.
 mixture results in good light-off and warmed-up performance for the
 conversion of both types of hydrocarbons. The presence of CeO₂ generally
 decreases light-off performance for most of these catalysts.

L2 ANSWER 3 OF 4 CAPLUS COPYRIGHT 2007 ACS on STN
 AN 1990:422873 CAPLUS
 DN 113:22873
 TI Homogeneous catalytic reduction of nitric oxide by olefins in the presence
 of palladium(II) chloride
 AU Cheng, C. H.; Sun, K. S.
 CS Dep. Chem., Natl. Tsing Hua Univ., Hsinchu, 30043, Taiwan
 SO Inorganic Chemistry (1990), 29(13), 2547-8
 CODEN: INOCAJ; ISSN: 0020-1669
 DT Journal
 LA English
 AB An aqueous solution of PdCl₂ and CuCl₂ catalyzes the reaction of NO with
 terminal
 olefins to give N₂O and the corresponding 2-ketone or aldehyde. The rate
 of catalysis at ambient temperature corresponds to 0.384, 0.255, 0.298, 0.190
 turnovers/h per Pd ion based on N₂O produced for ethylene,
 propylene, 1-butene, and 1-hexene, resp. PdCl₂ alone in a
 DMSO-H₂O (9:1, volume/volume) mixture is also an effective catalyst for the
 reduction of NO by terminal olefins. The observed catalysis is inhibited by
 Cl⁻
 and by H⁺.

L2 ANSWER 4 OF 4 CAPLUS COPYRIGHT 2007 ACS on STN
 AN 1985:112790 CAPLUS
 DN 102:112790
 TI Studies on the reactivity of allylic compounds catalyzed by a palladium[0]
 complex - the effect of leaving groups
 AU Huang, Jiling; Li, Bihua; Lu, Xiyan
 CS Shanghai Inst. Org. Chem., Acad. Sin., Shanghai, Peop. Rep. China
 SO Kexue Tongbao (Foreign Language Edition) (1984), 29(8), 1038-41
 CODEN: KHTPB; ISSN: 0454-0948
 DT Journal
 LA English
 AB The effect of leaving groups X were determined in the reaction of the
 RCH:CHCH₂X [X = OAc, O₂Cet, OP(O)(OCH₂CH:CH₂)₂, OPh, Br, NH₂, NHC(S)NH₂,
 etc.; R = e.g., H] with BH₄; catalyzed by Pd(PPh₃)₄, to give
 propylene. The best leaving group was OAc.

=> s palladium carbide
172232 PALLADIUM
219213 CARBIDE
L3 97 PALLADIUM CARBIDE
(PALLADIUM(W) CARBIDE)

=> s l3 and propylene
192172 PROPYLENE
L4 1 L3 AND PROPYLENE

=> d bib abs

L4 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2007 ACS on STN
AN 1998:352170 CAPLUS
DN 129:54753
TI Primary hydroxy-modified fluoropolyethers with narrow molecular weight
distribution and their manufacture
IN Tsuji, Makoto; Nanbu, Hiromi
PA Kao Corp., Japan
SO Jpn. Kokai Tokkyo Koho, 11 pp.
CODEN: JKXXAF
DT Patent
LA Japanese
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 10147639	A	19980602	JP 1996-310406	19961121
PRAI	JP 1996-310406		19961121		
OS	MARPAT 129:54753				

AB The fluoropolyethers represent HO(AO)nR1 [R1 = H, (un)substituted C1-30 hydrocarbyl, alkylsilyl, acyl; A = ethylene, propylene, CH2C(CF3)H; n = 5-2000] wherein A groups may be same or different and contain ≥5 CH2C(CF3)H. The fluoropolyethers are manufactured by (1) random or block polymerizing trifluoropropylene oxide (I) with ethylene oxide and/or propylene oxide in the presence of only R2OH metal alkoxides [R2 = CRaRbRc, p-RdC6H4CH2; Ra, Rb, Rc = C1-10 alkyl, (alkyl)phenyl, benzyl; Rd = H, C1-10 alkyl, alkoxy] or R2OH and alkali metal hydroxides or alkali metal alkoxides, (2) stopping the polymerization using terminators, and (3) decomposing initiation end groups with acids or reductants. Thus, I was polymerized in the presence of BuOH at ≤40° for 3 h, reacted with 1N HCl, and reacted with chlorosulfonic acid for 12 h to give 89% HO[CH2CH(CF3)O]nH (n = 12 on the average) showing Mw/Mn 1.1 and Mn 1.9 + 103.

=> s l3 and reduction
332043 REDUCTION
L5 2 L3 AND REDUCTION

=> d 1-2 bib abs

L5 ANSWER 1 OF 2 CAPLUS COPYRIGHT 2007 ACS on STN
AN 1995:1000392 CAPLUS
DN 124:100540
TI Effect of oxygen implantation on the electrochemical properties of
palladium
AU Fujihana, Takanobu; Ueshima, Masato; Takahashi, Katsuo; Iwaki, Masaya
CS Advanced Technology Inc., Kanagawa, 245, Japan
SO Materials Research Society Symposium Proceedings (1995), 354 (Beam-Solid Interactions for Materials Synthesis and Characterization), 287-92
CODEN: MRSPDH; ISSN: 0272-9172
PB Materials Research Society
DT Journal

LA English
 AB The electrochem. properties of O+-implanted Pd measured by cyclic voltammetry in a 0.25 mol dm⁻³ H₂SO₄ solution were studied in relation to their composition and structure. Implantation of 160+ was performed with doses between 1017 and 1018 ions cm⁻² at 150 keV, and at nearly room temperature SIMS; ERD combined with RBS, and XRD were used to analyze the composition depth profile and structure of O+-implanted Pd surface layers. The H atoms were accumulated with a Gaussian distribution and carbon materials containing the solid solution of PdCx (x = 0.13-0.15) were also formed in the near surface layers during O+-implantation. The distribution of implanted oxygen changed from Gaussian to trapezoidal as the dose increased, accompanied by the crystal growth of Pd(OH)₂, and simultaneously, the amount of accumulated H atoms increased. The voltammetric measurements revealed that with an increase in the dose, the hydrogen absorption was suppressed at the early stage of sweep cycles, and at the final stage, the redox reaction of both hydrogen and Pd was activated. Probably the carbon materials containing the PdCx formed during O+-implantation suppress the hydrogen absorption, and the metallic Pd like a Pd-black formed by the reduction of Pd(OH)₂ during voltammetric measurements causes the electrochem. activation of O+-implanted Pd.

L5 ANSWER 2 OF 2 CAPLUS COPYRIGHT 2007 ACS on STN

AN 1990:216185 CAPLUS

DN 112:216185

TI Rhodium(I)- or ruthenium(II)-catalyzed direct coupling of vinyl ketones with aldehydes and the subsequent reduction to give aldol derivatives anti-selectively

AU Sato, Susumu; Matsuda, Isamu; Shibata, Masahiro

CS Fac. Eng., Nagoya Univ., Nagoya, 464-01, Japan

SO Journal of Organometallic Chemistry (1989), 377(2-3), 347-56
 CODEN: JORCAI; ISSN: 0022-328X

DT Journal

LA English

OS CASREACT 112:216185

AB A vinyl ketone reacts with an aldehyde to give an α -methylene- β -hydroxyalkanone with the concomitant formation of the vinyl ketone dimer in the presence of catalytic amount of RhH(PPh₃)₄ or RuH₂(PPh₃)₄ under almost neutral conditions. The selectivity of the cross-coupling product is remarkably improved in the presence of an extra mole of aldehyde. This type of cross-coupling is explained by the intermediacy of the transition-metal enolate, which is formed by the Michael-type addition of M-H to a vinyl ketone. The subsequent hydrogenation of the C-C double bond of the α -methylene- β -hydroxyalkanone proceeds readily to give aldol derivs. in the presence of the catalyst. [Rh(COD)(DPPB)]PF₆, [COD = 1,5-cyclooctadiene, DPPB = 1,4-bis(diphenylphosphino)butane] is the best choice of catalyst and gives aldol derivs. anti-selectively.

=> fil reg

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	37.17	38.43
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	-5.46	-5.46

FILE 'REGISTRY' ENTERED AT 13:08:06 ON 10 NOV 2007

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2007 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 9 NOV 2007 HIGHEST RN 952797-35-2
DICTIONARY FILE UPDATES: 9 NOV 2007 HIGHEST RN 952797-35-2

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH June 29, 2007

Please note that search-term pricing does apply when
conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and
predicted properties as well as tags indicating availability of
experimental property data in the original document. For information
on property searching in REGISTRY, refer to:

<http://www.cas.org/support/stngen/stndoc/properties.html>

=> d his

(FILE 'HOME' ENTERED AT 13:00:20 ON 10 NOV 2007)

FILE 'CAPLUS' ENTERED AT 13:03:48 ON 10 NOV 2007

L1 728 S REDUCTION OF PALLADIUM
L2 4 S L1 AND PROPYLENE
L3 97 S PALLADIUM CARBIDE
L4 1 S L3 AND PROPYLENE
L5 2 S L3 AND REDUCTION

FILE 'REGISTRY' ENTERED AT 13:08:06 ON 10 NOV 2007

=> s pdc0.16/mf

L6 0 PDC0.16/MF

=> s pdc.16/mf

L7 0 PDC.16/MF

=> s pd1c.16/mf

L8 0 PD1C.16/MF

=> s pd1c.85/mf

L9 0 PD1C.85/MF